

武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

Recombinant Interleukin 22 Receptor Alpha 2 (IL22Ra2)

Catalog No.: TD-TP05648 50µg

Sequence Information

Species: Human Gene ID:116379

Swiss Prot:Q969J5 Synonyms:CRF2-S1; IL-22BP; ZcytoR16;

Cytokine receptor class-II member 10;

Cytokine receptor family type 2, soluble

1; Interleukin-22-binding protein

Residues:Thr22~Pro263

TQSTHESLKPQRVQFQSRNFHNILQWQPGRALTGNSSVYFVQYKIMFSCSMKSS

HQKPSGCWQHISCNFPGCRTLAKYGQRQWKNKEDCWGTQELSCDLTSETSDIQE

PYYGRVRAASAGSYSEWSMTPRFTPWWETKIDPPVMNITQVNGSLLVILHAPNL

PYRYQKEKNVSIEDYYELLYRVFIINNSLEKEQKVYEGAHRAVEIEALTPHSSY

CVVAEIYQPMLDRRSQRSEERCVEIP

Product Information

Source: Recombinant expression.

Host: E.coli

Tags: N-terminal His-Tag

Subcellular Location: Secreted.

Purity: >90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 400µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.0

Predicted Molecular Mass: 31.9kDa

Accurate Molecular Mass: 32kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

STORAGE AND STABILITY



武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. Gene Sequencing (Extract)




Figure 2. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.