

武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

### Eukaryotic Cluster Of Differentiation 56 (CD56)

Catalog No.: TD-TP04993 50µg

### **Sequence Information**

Species: Human Swiss Prot:P13591 Gene ID:4684 Synonyms:CD56; NCAM1; MSK39; N-CAM-1; NCAM; Neural Cell Adhesion Molecule

Residues:Leu20-Pro603

LQVDIVPSQGEISVGESKFFLCQVAGDAKDKDISWFSPNGEKLTPNQQRISVVW NDDSSSTLTIYNANIDDAGIYKCVVTGEDGSESEATVNVKIFQKLMFKNAPTPQ EFREGEDAVIVCDVVSSLPPTIIWKHKGRDVILKKDVRFIVLSNNYLQIRGIKK TDEGTYRCEGRILARGEINFKDIQVIVNVPPTIQARQNIVNATANLGQSVTLVC DAEGFPEPTMSWTKDGEQIEQEEDDEKYIFSDDSSQLTIKKVDKNDEAEYICIA ENKAGEQDATIHLKVFAKPKITYVENQTAMELEEQVTLTCEASGDPIPSITWRT STRNISSEEKASWTRPEKQETLDGHMVVRSHARVSSLTLKSIQYTDAGEYICTA SNTIGQDSQSMYLEVQYAPKLQGPVAVYTWEGNQVNITCEVFAYPSATISWFRD GQLLPSSNYSNIKIYNTPSASYLEVTPDSENDFGNYNCTAVNRIGQESLEFILV QADTPSSPSIDQVEPYSSTAQVQFDEPEATGGVPILKYKAEWRAVGEEVWHSKW YDAKEASMEGIVTIVGLKPETTYAVRLAALNGKGLGEISAASEF

### **Product Information**

Source: Eukaryotic expression.

Host: 293F cell

Tags: N-terminal His Tag and C-terminal Fc Region of Human

lgG1

Subcellular Location: Membrane, Secreted.

**Purity: >90%** 

Traits: Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

### Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 4.3

Predicted Molecular Mass: 95.8kDa

电话: 027-87561633 手机: 189-8623-2100 QQ客服: 121545059 网址:http://www.itiande-neuron.com 公司地址: 武汉东湖新技术开发区高新大道666号光谷生物城生物创新园B1栋b274-9室



武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

Accurate Molecular Mass: 100kDa as determined by SDS-PAGE reducing conditions.

# [ <u>USAGE</u> ]

Reconstitute in  $ddH_2O$  to a concentration of 0.1-0.5 mg/mL. Do not vortex.

# [ STORAGE AND STABILITY ]


Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]



### Figure 1. SDS-PAGE

## [<u>IMPORTANT NOTE</u>]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.