

### 武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

#### **Recombinant Cluster of Differentiation 59 (CD59)**

Catalog No.: TD-TP01524 50µg

**Sequence Information** 

Species: Human Gene ID:966

Swiss Prot:P13987 Synonyms:Protectin; MIC11; MIN1; MIN2;

MIN3; MSK21; MAC-IP; MACIF;

MIRL; HRF20; Complement Regulatory

Protein; 20 kDa homologous restriction

factor; Membrane attack complex

inhibition factor

Residues:Leu26~Asn102

LQCYNCPNPTADCKTAVNCSSDFDACLITKAGLQVYNKCWKFEHCNFNDVTTRL

RENELTYYCCKKDLCNFNEQLEN

**Product Information** 

Source: Prokaryotic expression.

Host: E. coli

Tags: N-terminal His Tag

Subcellular Location: Secreted, Extracellular matrix.

**Purity: >97%** 

Traits: Freeze-dried powder

**Buffer formulation:**PBS, pH7.4, containing 0.1% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.5

Predicted Molecular Mass: 15.7kDa

Accurate Molecular Mass: 19kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

## 武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ IDENTIFICATION ]




Figure 1. Gene Sequencing (Extract)




Figure 2. SDS-PAGE



# 武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.