

武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

Eukaryotic Cluster Of Differentiation 33 (CD33)

Catalog No.: TD-TP04909 50µg

Sequence Information

Species: Human Swiss Prot:P20138 Gene ID:945 Synonyms:SIGLEC3; Gp67; p67; Myeloid cell surface antigen CD33

Residues: Asp18-His259

PNFWLQVQESVTVQEGLCVLVPCTFFHPIPYYDKNSPVHGYWFREGAIISRDSP VATNKLDQEVQEETQGRFRLLGDPSRNNCSLSIVDARRRDNGSYFFRMERGSTK

YSYKSPQLSVHVTDLTHRPKILIPGTLEPGHSKNLTCSVSWACEQGTPPIFSWL

SAAPTSLGPRTTHSSVLIITPRPQDHGTNLTCQVKFAGAGVTTERTIQLNVTYV

PQNPTTGIFPGDGSGKQETRAGVVH

Product Information

Source: Eukaryotic expression. Host: 293F cell Tags: N-terminal His Tag Subcellular Location: Membrane. Purity: >95% Traits: Freeze-dried powder Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. Original Concentration: 800µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.0

Predicted Molecular Mass: 28.4kDa

Accurate Molecular Mass: 48kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in ddH_2O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

电话: 027-87561633 手机: 189-8623-2100 QQ客服: 121545059 网址:http://www.itiande-neuron.com 公司地址: 武汉东湖新技术开发区高新大道666号光谷生物城生物创新园B1栋b274-9室



Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

武汉天德生物科技有限公司 Wuhan Tanda Biotechnology Co., Ltd

[IDENTIFICATION]




Figure 1. SDS-PAGE

[<u>IMPORTANT NOTE</u>]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.